

Feedback-Bericht Disease Management Programm Koronare Herzkrankheit 2006/1

Literatur zum Zusatzbericht „Differenzialtherapie mit Thrombozyten-Aggregationshemmern in der Sekundärprävention der koronaren Herzkrankheit“

- 1: Siegel-Axel, D., Langer, H., Lindemann, S. & Gawaz, M. (2006). Die Rolle von Thrombozyten bei Entzündungs- und Atheroskleroseprozessen. *Medizinische Klinik*, 101 (6), 467–475.
- 2: Antithrombotic Trialists' Collaboration (2002). Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *British Medical Journal*, 324 (7329), 71–86.
- 3: Hamm, C. W., Arntz, H.-R., Bode, C., Giannitsis, E., Katus, H., Levenson, B., Nordt, T., Neumann, F. J., Tebbe, U., Zahn, R. (2004). Leitlinien der Deutschen Gesellschaft für Kardiologie. Akutes Koronarsyndrom (ACS). Teil 2: Akutes Koronarsyndrom mit ST-Hebung. *Zeitschrift für Kardiologie*, 93 (4), 324–341.
- 4: Chen, Z.M., Jiang, L. X., Chen, Y. P., Xie, J. X., Pan, H. C., Peto, R., Collins, R., Liu, L. S.; COMMIT Clopidogrel and Metoprolol in Myocardial Infarction Trial collaborative group (2005). Addition of clopidogrel to aspirin in 45852 patients with acute myocardial infarction: Randomised placebo-controlled trial. *Lancet*, 366 (9497), 1607–1621.
- 5: Sabatine, M. S., Cannon, C. P., Gibson, C. M., Lopez-Sendon, J. L., Montalescot, G., Theroux, P., Lewis, B. S., Murphy, S. A., McCabe, C. H., Braunwald, E.; CLARITY-TIMI 28 Investigators (2005). Effect of clopidogrel pretreatment before percutaneous coronary intervention in patients with ST-elevation myocardial infarction treated with fibrinolytics: The PCI-CLARITY study. *Journal of the American Medical Association*, 294 (10), 1224–1232.
- 6: Sabatine, M. S., Cannon, C. P., Gibson, C. M., Lopez-Sendon, J. L., Montalescot, G., Theroux, P., Claeys, M. J., Cools, F., Hill, K. A., Skene, A. M., McCabe, C. H., Braunwald, E.; CLARITY-TIMI 28 Investigators (2005). Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *New England Journal of Medicine*, 352 (12), 1179–1189.
- 7: Silber, S., Albertsson, P., Avilés, F. F., Camici, P. G., Colombo, A., Hamm, C., Jørgensen, E., Marco, J., Nordrehaug, J. E., Ruzyllo, W., Urban, P., Stone, G. W., Wijns, W.; Task Force for Percutaneous Coronary Interventions of the European Society of Cardiology (2005). Guidelines for percutaneous coronary interventions. *European Heart Journal*, 2005, 26 (8), 804–847.
- 8: CAPRIE Steering Committee (1996). A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). *Lancet*, 348 (9038), 1329–1339.
- 9: Bhatt, D. L., Fox, K. A. A., Hacke, W., Berger, P. B., Black, H. R., Boden, W. E., Cacoub, P., Cohen, E. A., Creager, M. A., Easton, J. D., Flather, M. D., Haffner, S. M., Hamm, C. W., Hankey, G. J., Johnston, S. C., Mak, K.-H., Mas, J.-L., Montalescot, G., Pearson, T. A., Steg, P. G., Steinhubl, S. R., Weber, M. A., Brennan, D. M., Fabry-Ribaudo, L., Booth, J., Topol, E. J., CHARISMA Investigators (2006). Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. *New England Journal of Medicine*, 354 (16), 1706–1717.
- 10: Antiplatelet Trialists' Collaboration (1994). Collaborative overview of randomised trials of antiplatelet therapy – I: Prevention of death, myocardial infarction, and stroke by prolonged antiplatelet therapy in various categories of patients. *British Medical Journal*, 308 (6921), 81–106.
- 11: McAnally, L. E., Corn, C. R., Hamilton, S. F. (1992). Aspirin for the prevention of vascular death in women. *Annals of Pharmacotherapy*, 26 (12), 1530–1534.
- 12: Hammes, H.-P., Bertram, B., Bornfeld, N., Danne, T., Kroll, P., Lemmen, K. D.; Scherbaum, W. A., Kiess, W., Deutsche Diabetes-Gesellschaft (Hg) (2004). *Diagnostik, Therapie und Verlaufskontrolle der diabetischen Retinopathie und Makulopathie. Evidenzbasierte Leitlinie DDG*. Düsseldorf: Deutsche Diabetes-Gesellschaft.
- 13: Steinhubl, S. R., Berger, P. B., Mann, J. T. 3rd, Fry, E. T., DeLago, A., Wilmer, C., Topol, E. J.; CREDO Investigators (2002). Early and sustained dual oral antiplatelet therapy following percutaneous coronary intervention: A randomized controlled trial. *Journal of the American Medical Association*, 288 (19), 2411–2420.
- 14: Mehta, S. R., Yusuf, S., Peters, R. J., Bertrand, M. E., Lewis, B. S., Natarajan, M. K., Malmberg, K., Rupprecht, H., Zhao, F., Chrolavicius, S., Copland, I., Fox, K. A.; Clopidogrel in Unstable angina to prevent Recurrent Events trial (CURE) Investigators (2001). Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: The PCI-CURE study. *Lancet*, 358 (9281), 527–533.
- 15: Mangano, D. T.; Multicenter Study of Perioperative Ischemia Research Group (2002). Aspirin and mortality from coronary bypass surgery. *New England Journal of Medicine*, 347 (17), 1309–1317.